

Future ocean acidification will be amplified by hypoxia in coastal habitats

Frank Melzner · Jörn Thomsen · Wolfgang Koeve ·
Andreas Oschlies · Magdalena A. Gutowska ·
Hermann W. Bange · Hans Peter Hansen · Arne Körtzinger

Received: 31 December 2011 / Accepted: 30 April 2012
© Springer-Verlag 2012

Abstract Ocean acidification is elicited by anthropogenic carbon dioxide emissions and resulting oceanic uptake of excess CO₂ and might constitute an abiotic stressor powerful enough to alter marine ecosystem structures. For surface waters in gas-exchange equilibrium with the atmosphere, models suggest increases in CO₂ partial pressure ($p\text{CO}_2$) from current values of ca. 390 μatm to ca. 700–1,000 μatm by the end of the century. However, in typically unequilibrated coastal hypoxic regions, much higher $p\text{CO}_2$ values can be expected, as heterotrophic degradation of organic material is necessarily related to the production of CO₂ (i.e., dissolved inorganic carbon). Here, we provide data and estimates that, even under current conditions, maximum $p\text{CO}_2$ values of 1,700–3,200 μatm can easily be reached when all oxygen is consumed at salinities between 35 and 20, respectively. Due to the nonlinear nature of the carbonate system, the approximate doubling of seawater $p\text{CO}_2$ in surface waters due to ocean acidification will most strongly affect coastal hypoxic

zones as $p\text{CO}_2$ during hypoxia will increase proportionally: we calculate maximum $p\text{CO}_2$ values of ca. 4,500 μatm at a salinity of 20 ($T = 10^\circ\text{C}$) and ca. 3,400 μatm at a salinity of 35 ($T = 10^\circ\text{C}$) when all oxygen is consumed. Upwelling processes can bring these CO₂-enriched waters in contact with shallow water ecosystems and may then affect species performance there as well. We conclude that (1) combined stressor experiments ($p\text{CO}_2$ and $p\text{O}_2$) are largely missing at the moment and that (2) coastal ocean acidification experimental designs need to be closely adjusted to carbonate system variability within the specific habitat. In general, the worldwide spread of coastal hypoxic zones also simultaneously is a spread of CO₂-enriched zones. The magnitude of expected changes in $p\text{CO}_2$ in these regions indicates that coastal systems may be more endangered by future global climate change than previously thought.

Communicated by S. Dupont.

F. Melzner (✉) · J. Thomsen
Marine Ecology, Helmholtz Centre for Ocean Research
(GEOMAR), Kiel, Germany
e-mail: fmelzner@ifm-geomar.de

W. Koeve · A. Oschlies
Biogeochemical Modelling, Helmholtz Centre for
Ocean Research (GEOMAR), Kiel, Germany

M. A. Gutowska
Biological Oceanography, Helmholtz Centre for
Ocean Research (GEOMAR), Kiel, Germany


H. W. Bange · H. P. Hansen · A. Körtzinger
Chemical Oceanography, Helmholtz Centre for
Ocean Research (GEOMAR), Kiel, Germany

Introduction

Currently, large research efforts are directed toward studying anthropogenic climate change effects on marine ecosystems. Global warming, primarily caused by increasing CO₂ emissions, has the potential to restructure marine ecosystems (IPCC 2007). In highly productive coastal ecosystems, other abiotic stressors play a role as well: eutrophication largely following the human footprint has led to a spread of low-oxygen regimes where benthic metazoan communities cannot survive (Diaz and Rosenberg 2008). While the biological effects of hypoxia and anoxia on marine metazoan heterotrophic organisms are fairly well understood, it has been sparsely acknowledged that hypoxia, a consequence of respiratory gas exchange, is necessarily coupled to an increase in total dissolved

inorganic carbon and CO_2 partial pressure ($p\text{CO}_2$). In this paper, we will give estimates of $p\text{CO}_2$ values in coastal marine and estuarine habitats now and within the next 50–100 years. To do so, we will calculate how respiration in hypoxic zones interacts with the invasion of anthropogenic CO_2 into the ocean (ocean acidification). Our estimates have important repercussions for the design of experimental studies currently underway to assess the effect of ocean acidification on marine biota. We further suggest that carbonate chemistry of coastal zones may be much more sensitive with respect to future global change than previously estimated. We finally try to point the reader's attention toward key physiological systems and mechanisms that might play a role in determining marine animal vulnerability in response to the hypoxia and ocean acidification (hypercapnia) stress complex.

Anthropogenic CO_2 emissions not only are leading to global warming but also change ocean chemistry. When CO_2 diffuses into seawater, it reacts with water and carbonic acid is formed. Carbonic acid rapidly dissociates into bicarbonate (HCO_3^-) and a proton and can further dissociate into the carbonate ion (CO_3^{2-}) and another proton:



The equilibrium constants of the inorganic carbonate system are such that under current abiotic conditions of temperature, salinity and alkalinity, roughly 90 % of the CO_2 is in the form of HCO_3^- and ~10 % in the form of CO_3^{2-} , while less than 1 % is dissolved CO_2 and carbonic acid (termed CO_2^*). The sum of the concentrations of the carbonate system species is termed total dissolved inorganic carbon (C_T):

$$C_T = [\text{CO}_2^*] + [\text{HCO}_3^-] + [\text{CO}_3^{2-}] \quad (2)$$

Any invasion of CO_2 from the atmosphere to the surface ocean leads to an increase in C_T and a shift in carbonate

system speciation, namely an increase in $[\text{HCO}_3^-]$, an increase in $p\text{CO}_2$, a decrease in $[\text{CO}_3^{2-}]$ and a decrease in pH (Brewer 1997, see Table 1). Currently (end of 2011), atmospheric $p\text{CO}_2$ is at 391 ppm and increases at a rate of 2 ppm year⁻¹ (NOAA/ESRL, www.esrl.noaa.gov/gmd/ccgg/trends, Cao and Caldeira 2008). It has been estimated that atmospheric $p\text{CO}_2$ will rise to values of 700–1,000 ppm by the end of the century (IPCC 2007). It is well documented that the past time change in atmospheric $p\text{CO}_2$ has led to an oceanic uptake of CO_2 and associated ocean acidification (Kleypas et al. 1999). With this continuing in future, we can expect a moderate increase in $[\text{HCO}_3^-]$, while the carbonate ion concentration $[\text{CO}_3^{2-}]$ decreases starkly (by ~40 %) and pH decreases by ~0.25 units by 2100 (Table 1). Decreases in $[\text{CO}_3^{2-}]$ lead to a reduction in the calcium carbonate saturation state (Ω) for the two common polymorphs, aragonite and calcite, that marine organisms use to construct their CaCO_3 shells or tests:

$$\Omega = [\text{CO}_3^{2-}] [\text{Ca}^{2+}] K_{\text{sp}}^{*-1} \quad (3)$$

where K_{sp}^* = apparent stoichiometric solubility coefficient for aragonite or calcite.

In the contemporary ocean surface waters, Ω typically is >1 for both aragonite and calcite (e.g., Cao and Caldeira 2008, see Table 1). Once Ω drops <1, dissolution of CaCO_3 is thermodynamically favored. The carbonate ion concentration, and hence Ω , is also a function of salinity, with lower values in brackish waters (Table 1).

Coastal hypoxia is a problem of global dimensions and has been a research focus during the last decades. Diaz and Rosenberg (2008) describe severely hypoxic zones from 400 systems worldwide, covering an area of >245,000 km². An oxygen concentration of about 60–70 μM (~20–30 % air saturation) has been suggested as an important hypoxic threshold value (e.g., Diaz and Rosenberg 2008; Vaquer-Sunyer and Duarte 2008; Levin et al. 2009), as pronounced sublethal effects (e.g., aberrant behavior, growth

Table 1 Carbonate system speciation ($S = 20$ vs. $S = 35$) in fully air-saturated seawater for scenarios $1\times$, $2\times$ and $3\times$ CO_2 , assuming constant total alkalinities (A_T)

Scenario	$p\text{CO}_2$ (μatm)	$C_{T,\text{fut}}$ ($\mu\text{mol kg}^{-1}$)	A_T ($\mu\text{mol kg}^{-1}$)	pH_{NBS}	Ω_{calc}	Ω_{arag}
$S = 20$, $[\text{O}_2] = 310.4 \mu\text{M}$						
$1\times \text{CO}_2$	390	1,877	1,976	8.18	2.22	1.33
$2\times \text{CO}_2$	780	1,948	1,976	7.90	1.23	0.74
$3\times \text{CO}_2$	1,170	1,987	1,976	7.73	0.85	0.51
$S = 35$, $[\text{O}_2] = 282 \mu\text{M}$						
$1\times \text{CO}_2$	390	2,153	2,355.5	8.17	3.57	2.27
$2\times \text{CO}_2$	780	2,262	2,355.5	7.90	2.07	1.32
$3\times \text{CO}_2$	1,170	2,317	2,355.5	7.74	1.47	0.93

A_T was calculated using an A_T versus salinity relationship for the Western Baltic Sea (Beldowski et al. 2010). All scenarios were calculated from A_T and $p\text{CO}_2$ using CO2SYS (Lewis and Wallace 1998)

reductions, alterations in reproductive effort) often start to be apparent in many coastal invertebrates under such conditions. Many seasonal hypoxic coastal systems ('dead zones' sensu Diaz and Rosenberg 2008) are mainly located along Northern European seas and estuaries, the North American east coast and the Gulf of Mexico, as well as the coasts of Japan and China, that is, these systems follow the patterns of the human footprint. Hypoxia and anoxia have profound effects on benthic ecosystems, resulting in habitat compression, loss of fauna and biodiversity and large decreases in secondary production (see Diaz and Rosenberg 2008). Generally, eutrophication is regarded as the primary causative agent for many coastal hypoxic systems. Elevated nutrient concentrations and dissolved organic matter (DOM), typically introduced into coastal areas by means of river discharge but also by land run off, lead to enhanced rates of production of particulate organic matter (POM), partly sinking out from the surface layer to deeper waters and the benthos. The subsequent bacterial degradation of organic matter depletes the oxygen content of the water column (see Gray et al. 2002). High degrees of stratification in temperate waters and associated suppression of ventilation from above make the subsurface waters ideal nurseries for hypoxia and anoxia in the summer and autumn months.

Except for a few exceptions (e.g., Truchot and Duhameljoue 1980; Burnett 1997; Seibel and Walsh 2003; Brewer and Peltzer 2009), the biological review literature has often neglected the fact that coastal hypoxia is necessarily coupled to acidification and that the degree of acidification in hypoxic regions may represent a significant additional abiotic stressor and, occasionally, an adaptive physiological effector. Heterotrophic degradation of organic matter always results in the production of carbon dioxide and therefore acidification. There are some good examples available of simultaneous measurements of $[O_2]$ and carbonate system speciation in hypoxic coastal regions. A linear relationship between pH and $[O_2]$ has been obtained for the Charleston Harbor Estuary (east coast USA, Burnett 1997), with minimum pH_{NBS} values of <7.0 and maximum pCO_2 values in excess of $10,000 \mu atm$ (Cochran and Burnett 1996). Inverse relationships between $\Delta[O_2]$ and ΔC_T have been documented for, for example, Tampa Bay (Florida, USA), where strong diurnal patterns exist due to daytime photosynthesis versus nighttime respiration (Yates et al. 2007), the Scheldt Estuary (Belgium), where almost equimolar changes in $[O_2]$ and C_T led to extremely high pCO_2 values of $>5,000 \mu atm$ (Frankignoulle et al. 1996), oxygen-minimum zones (Paulmier et al. 2011), the very large Gulf of Mexico and East China Sea hypoxic zones (Cai et al. 2011), and urban estuaries along the western Pacific coast (Feely et al. 2010).

In the following, we present data from a 10-year data set on $[O_2]$ and pH_{NBS} from the Boknis Eck Time Series Station (BE) in the Western Baltic Sea to illustrate the close relationship between both parameters in a seasonal hypoxic coastal system and calculate the carbonate system speciation at BE using C_T and A_T measurements obtained in 2009–2010. We then present calculations of pCO_2 in coastal hypoxic habitats and try to extrapolate pCO_2 values for such systems, assuming a doubling (tripling) in surface seawater pCO_2 during this century (the next centuries), with particular reference to the high sensitivity of estuarine systems. We discuss the implications of these results for ongoing efforts in hypoxia and ocean acidification research.


Methods

Time series data

Time series data from the Boknis Eck Time Series Station (BE) in Eckernförde Bay (greater Kiel Bay, $54^\circ 31' 77'' N$, $10^\circ 02' 36'' E$, see Fig. 1) were collected on monthly cruises as described in Hansen et al. (1999). Salinity, temperature and pH_{NBS} were measured with sensors attached to a CTD rosette, and oxygen concentrations were measured using the Winkler method (Grasshoff et al. 1983). The pH electrode was calibrated before use using pH buffers (NBS) prepared according to Grasshoff et al. (1983). Water samples for $[O_2]$ analysis were obtained from discrete depths (1, 5, 10, 15, 20, 25, 27 m). For our analysis, data were filtered for records 20 m and deeper with simultaneous measurements of oxygen, salinity, temperature and pH_{NBS} . This resulted in a data set of 280 complete records, with between 18 and 29 samples per month and spanning a time interval of 10 years between 1986 and 1995. In order to calculate seawater pCO_2 during this time interval, we estimated total alkalinity using the salinity– A_T relationship obtained by Beldowski et al. (2010) for the Western Baltic Sea. The carbonate system speciation was then calculated from estimated A_T and measured pH_{NBS} using CO2SYS (Lewis and Wallace 1998) with $KHSO_4$ dissociation constants according to Dickson (1990) and dissociation constants K_1 and K_2 according to Roy et al. (1993).

In order to obtain more precise information on seasonal changes in pCO_2 at station BE, we sampled water for accurate determinations of A_T and C_T during an entire seasonal cycle. Water samples were taken from 20 and 25 m depths on monthly cruises with the research vessel FK Littorina between January 2009 and March 2010 in the framework of BE time series measurements. Samples were stored in 500-ml Duran glass bottles, immediately poisoned using 100 μl saturated mercuric chloride solution and

Fig. 1 Map of Kiel Bay in the Western Baltic Sea indicating the location of Boknis Eck Time Series Station (BE) on the entrance of Eckernförde Bay ($54^{\circ}31'N$, $10^{\circ}02'E$, max. water depth 28 m)



stored at room temperature until analysis. C_T and A_T were measured by coulometric and potentiometric titration using SOMMA and VINDTA autoanalyzers, respectively (Minrop et al. 2000; Dickson et al. 2007). A_T and C_T measurements were corrected using measurements of certified seawater standards (Dickson et al. 2007). pCO_2 and pH_{NBS} were then calculated using the CO2SYS program as described above. All other parameters (S , T , O_2) were determined as described by Hansen et al. (1999).

Surface seawater pH_{NBS} was measured on September 6, 2009, between 11:00 and 20:00 around Kiel Fjord following 2 days of strong south–westerly winds that caused upwelling of deeper water layers. Measurements were taken manually in water depths of ca. 0.5 m using a WTW pH meter equipped with a Sentix 81 pH_{NBS} electrode. More data on surface pH variability and carbonate chemistry of Kiel Fjord in 2008–2010 are given in Thomsen et al. (2010).

Carbonate system calculations

The carbonate system speciation in Table 1 and Figs. 2 and 3 was calculated using CO2SYS (Lewis and Wallace 1998), with K_1 and K_2 from Roy et al. (1993), assuming silicate and phosphate = $0 \mu\text{mol kg}^{-1}$ for present and future seawater pCO_2 values ($1 \times CO_2 = 390 \mu\text{atm}$, $2 \times CO_2 = 780 \mu\text{atm}$, $3 \times CO_2 = 1,170 \mu\text{atm}$) and total alkalinities (A_T) for salinities of 20–35 based on an A_T salinity

relationship obtained by Beldowski et al. (2010). Additional generation of C_T from respiration ($C_{T,resp}$) in hypoxic regions was estimated using a $\Delta[O_2]/\Delta C_T$ conversion factor of -1.34 from Körtzinger et al. (2001; this factor is equivalent to a respiratory quotient (RQ) of 0.75) and an initial $[O_2]$ of $310.4 \mu\text{M}$ (at $S = 20$, $T = 10^\circ\text{C}$) and $282 \mu\text{M}$ (at $S = 35$, $T = 10^\circ\text{C}$), assuming 100 % air saturation. Future C_T ($C_{T,fut}$) thus was calculated as the sum of current C_T ($C_{T,curr}$), C_T from respiration processes ($C_{T,resp}$) and C_T from the invasion of anthropogenic CO_2 , that is, ocean acidification ($C_{T,oa}$):

$$C_{T,fut} = C_{T,curr} + \Delta C_{T,oa} + \Delta C_{T,resp} \quad (4)$$

Seawater pCO_2 , pH_{NBS} and Ω_{arag} values for the hypoxic model systems in Fig. 5 were then calculated from $C_{T,fut}$ and A_T . A_T remains constant when additional CO_2 is introduced into the system (e.g., Dickson et al. 2007). Our calculations do not include potential effects of future temperature change (which would lower oxygen solubility but increase the rate of respiratory decline in $[O_2]$, thus further increasing duration and severity of the hypoxic period). We also did not consider potential changes in A_T in a fully anoxic system. Thus, the calculations presented in Fig. 5 should be viewed as a rough indicator of magnitudes of change in pCO_2 that can be expected in coastal hypoxic systems.




Fig. 2 Seasonal variations in salinity, temperature, [O₂], pH_{NBS} (measured), pCO₂ and Ω_{arag} (estimated) between 1986 and 1995 at Boknis Eck Time Series Station (BE). Open symbols in **a**, **b** indicate individual measurements, filled symbols are monthly mean values, and error bars are standard deviations. Carbonate system calculations

(pCO₂, Ω_{arag}) are based on measured pH_{NBS} and calculated A_T using an A_T-salinity relationship for the Western Baltic Sea (Beldowski et al. 2010). [O₂], salinity and temperature at BETS have previously been published by Hansen et al. (1999) for the period 1957–1997




Fig. 3 Seasonal variation in carbonate chemistry at Boknis Eck Time Series Station (BE) between 2009 and 2010 in relation to [O₂]. Linear (a, b) and nonlinear (c) regressions of pH_{NBS}, dissolved inorganic

carbon (C_T) and pCO₂ versus seawater [O₂]. pH_{NBS} and pCO₂ were calculated from A_T and C_T measurements. Measurements were taken at the same location as those represented in Fig. 2

Results

Time series data

At the Boknis Eck Time Series (BE) Station in Kiel Bay (Fig. 1), Western Baltic Sea, stratification during the summer months results in the formation of a deeper water body at depths greater than 15 m that is shielded by a thermohaline pycnocline and therefore cannot equilibrate with the atmosphere until storms aid convective mixing in late autumn. From March–April onwards, [O₂] in the bottom layers (>20 m) decreases linearly due to respiratory processes at a rate of 1.5–2 μM day⁻¹, typically reaching values of 0–50 μM in August and September (Hansen et al. 1999). [O₂] and pH_{NBS} were measured simultaneously at the BE Time Series Station between 1986 and 1995 (Hansen et al. 1999). Monthly averages of pH_{NBS} followed the pattern in [O₂] (Fig. 2c, d), with an average minimum pH_{NBS} of <7.5 observed in August and September. Calculating seawater pCO₂ and Ω_{arag} from pH_{NBS} and (estimated) A_T illustrates that most likely, average pCO₂ >1,000 μatm was encountered during June–November and pCO₂ >2,000 μatm during August and September (Fig. 2e). Peak pCO₂ values exceeded 3,000 μatm in autumn. Average Ω_{arag} mirrored pCO₂, reaching average minimum values of <0.5 during June–September and never

rising above 1 during the entire seasonal cycle. Our accurate carbonate system determinations at BE Time Series Station during 2009–2010 obtained from the same depth horizon (Fig. 3) essentially confirm the rough estimates of pCO₂ obtained using the 1986–1995 data set: seawater pH_{NBS} declined linearly (Fig. 3a), C_T increased linearly (Fig. 3b) and pCO₂ increased exponentially (Fig. 3c) with declining [O₂]. Peak pCO₂ values of 2,500–3,500 μatm were observed when [O₂] fell below 50 μM. While it could be argued that such stressful conditions are only encountered by benthic communities below thermohaline pycnoclines, upwelling processes can bring shallow coastal ecosystems in contact with the ‘corrosive’ subsurface waters (sensu Feely et al. 2008) that contain large amounts of respiratory CO₂. Figure 4 gives such an example: surface water pH_{NBS} values <7.5 were measured in September 2009 in the inner regions of Kiel Fjord following 2 days of strong southwesterly winds, while the outer parts of the Fjord retained a pH_{NBS} of 7.9–8.1. Due to the long chemical equilibration time scale, upwelled water retains high pCO₂ for long times, while physically dissolved oxygen equilibrates more rapidly with the atmosphere. Thomsen et al. (2010) demonstrated that surface water (<1 m depth) pCO₂ in Kiel Fjord follows the seasonal cycle observed in deeper layers of Kiel Bay (Fig. 2), albeit with a higher variability and lower pCO₂.




Fig. 4 Upwelling of low-pH water in Kiel Fjord. Surface seawater pH_{NBS} measured on September 6, 2009, between 11:00 and 20:00 around Kiel Fjord in water depths of ca. 0.5 m

Model calculations

Figure 5a–c gives a simulation of the changes in carbonate system speciation that can be expected during a typical seasonal hypoxic event for a salinity of 20 at a temperature of 10 °C for current pCO_2 ($1 \times CO_2 = 390 \mu atm$) and two future scenarios ($2 \times CO_2 = 780 \mu atm$, $3 \times CO_2 = 1,170 \mu atm$; Table 1). Under current conditions ($1 \times CO_2$), patterns calculated resemble the measurements obtained at BE time series during 2009–2010 (Fig. 3), with pCO_2 increasing exponentially (Fig. 5a), pH decreasing linearly (Fig. 5b) and Ω_{arag} decreasing exponentially (Fig. 5c) with decreasing $[O_2]$. Maximum pCO_2 (ca. 3,200 μatm), minimum pH_{NBS} (ca. 7.3) and Ω_{arag} (ca. 0.2) at $[O_2] = 0 \mu M$ are close to values observed or estimated for the field at Time Series Station BE (Figs. 2, 3). Increased C_T in the future ($C_{T, fut}$) due to ocean acidification ($\Delta C_{T, OA}$) leads to

an enhanced sensitivity of the system during hypoxia: a doubling of surface pCO_2 to 780 μatm results in an increase in pCO_2 to ca. 4,500 μatm when all oxygen is consumed, thus a relative increase by ca. 1,300 μatm in comparison with the $1 \times CO_2$ scenario. The $3 \times CO_2$ scenario would lead to maximum pCO_2 values in excess of 5,000 μatm at $[O_2] = 0 \mu M$. As pCO_2 is the primary driver affecting animal biology, these large changes are of particular relevance. Correspondingly, Ω_{arag} is affected very strongly by $2 \times$ and $3 \times CO_2$ scenarios, with undersaturation occurring at full air saturation already and very low Ω_{arag} (<0.5) being reached at relatively high oxygen content of the seawater (250–300 μM). Minimum pH_{NBS} values between 7.1 and 7.2 are reached in $2 \times$ and $3 \times CO_2$ scenarios when all oxygen is consumed.

Figure 5d–f gives a simulation of the effects of salinity on carbonate system speciation when $[O_2] = 0 \mu M$. The figures clearly illustrate that increases in pCO_2 are more severe in brackish scenarios, which is related to the lower A_T and hence buffer capacity. Highest pCO_2 values of ca. 3,200 μatm at $S = 20$ contrast with maximum pCO_2 values of ca. 1,700 μatm at $S = 35$ ($1 \times CO_2$). Future scenarios indicate that the influence of salinity on maximum pCO_2 decreases, which leads to relatively higher increases in pCO_2 in the high-salinity scenarios, with maximum pCO_2 values ca. 4,500 μatm at $3 \times CO_2$ and $S = 35$ (Fig. 5d). Similarly, the relative sensitivity of seawater pH_{NBS} and Ω_{arag} to salinity changes decreases from $1 \times$ to $2 \times$ and $3 \times CO_2$ scenarios (Fig. 5e, f).


Discussion

Our field data and simulations indicate exponential increases in pCO_2 with declining $[O_2]$ and a very high sensitivity, particularly of estuarine habitats, to the combined impacts of C_T increases due to ocean acidification and due to respiration. Magnitudes of pCO_2 reached now and in future scenarios in such systems have been shown to strongly impact animal biology. In the following sections, we will briefly discuss biological impacts of ocean acidification (hypercapnia) and of hypoxia, as well as of those of hypoxic hypercapnia, on animal physiology and fitness. We argue that it is important in experimental research to consider both factors simultaneously, as both probably significantly impact the organisms' energy budgets and, ultimately, fitness.

Ocean acidification (hypercapnia)

Biological effects of ocean acidification on heterotrophic metazoans are not understood well at present, but it is clear that at least three effects play a role: (1) direct effects of

Fig. 5 Seawater $p\text{CO}_2$, pH_{NBS} and Ω_{arag} calculated for $S = 20$, $T = 10^\circ\text{C}$ in response to changes in $[\text{O}_2]$ (**a, b, c**) and in response to salinity changes between $S = 20$ – 35 , $T = 10^\circ\text{C}$ for $[\text{O}_2] = 0\ \mu\text{M}$ (**d, e, f**). All calculations were made using CO2SYS software (Lewis and Wallace 1998). Thin line: $1\times\text{CO}_2$, intermediately thick line: $2\times\text{CO}_2$, thick line: $3\times\text{CO}_2$



altered carbonate chemistry on external shell dissolution in calcifiers, (2) pathological effects of an altered acid–base status on biomineralization and physiological processes and (3) impacts of an altered acid–base status on species’ energy budgets. (1) Direct effects of low $[\text{CO}_3^{2-}]$ have been observed in some invertebrates: shell dissolution was observed at $\Omega < 1$ in pelagic snails with only thin (or no) organic coating protecting their shells (pteropods, e.g., Lischka et al. 2011). However, recent studies indicate that net calcification of exoskeletons proceeds at seawater $\Omega \ll 1$ in many marine metazoans (e.g., Ries et al. 2009; Thomsen et al. 2010; Dupont et al. 2010), even in those that experience external shell dissolution at the same time

(Lischka et al. 2011). The organic matrix that is a crucial component of all skeletons and tests of marine organisms, as well as organic shell covers, probably is the chief reason that prevents external dissolution at $\Omega < 1$ (see Ries et al. 2009; Tunnicliffe et al. 2009; Thomsen et al. 2010). (2) Indirect effects of ocean acidification can result from disturbances in extra- and intracellular acid–base balance. All heterotrophic metazoans studied maintain positive diffusion gradients of CO_2 toward the surrounding medium, resulting in relatively high body fluid (extracellular) $p\text{CO}_2$ values of ca. 1,000–4,000 μatm (Melzner et al. 2009, Fig. 6). Thus, $p\text{CO}_2$ is the main effect or that impacts heterotrophic metazoans during ocean acidification as it




Fig. 6 Schematic illustration depicting the effects of seawater $p\text{CO}_2$ on extracellular (blood, hemolymph or coelomic fluid) $p\text{CO}_2$ in marine animals: typically, a gradient in $p\text{CO}_2$ of 1,000–4,000 μatm (here: 2,000 μatm) is maintained between extracellular fluids and seawater in order to drive diffusive excretion of metabolic CO_2 . It is evident from this diagram that the exponential increases in $p\text{CO}_2$ that go along with hypoxic hypercapnia lead to extreme extracellular acid–base disturbance. Increased extracellular $p\text{CO}_2$ will either result in very high extracellular $[\text{HCO}_3^-]$ in extracellular pH-regulating animals or in very low pH and Ω_{arag} in animals that do not regulate extracellular pH. Both scenarios can lead to pathologies and disturbances of the energy budget (see text)

results in elevated body fluid $p\text{CO}_2$ so respiratory diffusion gradients can be maintained. Some organisms compensate this disturbance by actively altering their extracellular carbonate system speciation (i.e., net accumulation of HCO_3^-) to stabilize extracellular pH (e.g., fish), whereas others do not (e.g., bivalves; see Seibel and Walsh 2003; Melzner et al. 2009 for review). Either way, CO_2 -induced changes in extracellular acid–base equilibria can have repercussions on functional processes: an altered blood carbonate system speciation may be responsible for pathologies such as hypercalcification observed in cephalopods, crustaceans and teleost fish (Gutowska et al. 2010) and has been demonstrated to lead to pathological behavioral disturbances in teleost fish (Munday et al. 2009; Dixon et al. 2010; Nilsson et al. 2012). Very high extracellular $p\text{CO}_2$ in animals that do not regulate extracellular pH can also lead to internal shell dissolution (Melzner et al. 2011) and might induce oxidative stress (Tomanek et al. 2011). (3) On the other hand, an unaltered carbonate system speciation may induce metabolic depression (Pörtner et al. 2005) or increase the energetic costs for homeostasis and calcification (Wood et al. 2008; Thomsen and Melzner 2010; Stumpp et al. 2011; Deigweier et al. 2009; Widdicombe and Spicer 2008). In many calcifying marine

animals, the initial step of calcification probably is the intracellular formation of an amorphous calcium carbonate (ACC) precursor phase (see Weiner and Addadi 2011 for a review). Thus, cell pH and ion homeostasis are tightly linked to biomineralization. As intracellular pH is maintained relatively constant in most marine animals even under hypercapnic conditions (e.g., Zange et al. 1990; Ellington 1993), hypercapnia-induced changes in extracellular $p\text{CO}_2$ and the dependent reductions in pH in weak pH regulators can adversely affect the energetics of ion exchange between intra- and extracellular compartments, primarily by increasing the driving force for protons to enter the cytosol. Thus, changes in seawater $p\text{CO}_2$ can potentially impact a range of cellular processes that translate into elevated energetic costs and cause energy budget reallocation (e.g., Dupont et al. 2010; Stumpp et al. 2011).

Hypoxia

Several recent contributions describe responses of marine organisms and communities to low oxygen concentrations (e.g., Grieshaber et al. 1994; Diaz and Rosenberg 1995, 2008; Burnett 1997; Boutilier 2001; Gray et al. 2002; Rabalais et al. 2002; Levin et al. 2009; Seibel 2011). While it is evident that common definitions on the use of units and sensitivity thresholds are missing in the literature (however, see Hofmann et al. 2011), two principle types of responses of heterotrophic marine metazoans with respect to declining oxygen concentrations can be distinguished: oxyregulators maintain rates of oxygen consumption with declining water $p\text{O}_2$, typically by increasing diffusion gradients across their gas-exchange epithelia. This can be achieved via (energy intensive) increases in ventilatory and/or circulatory convection effort, increases in perfused gas-exchange areas and decreases in diffusion distances (e.g., see Grieshaber et al. 1994; Burnett 1997; Farrell and Richards 2009; Richards 2009 for biochemical and molecular adaptation mechanisms). However, once a critical oxygen partial pressure ($p\text{O}_{2,\text{crit}}$) is reached, oxygen consumption and demand are downregulated. Eventually, anaerobic metabolism is needed to support the organism's energetic requirements (see Pörtner and Grieshaber 1993; Grieshaber et al. 1994). Oxyconforming animals do not maintain oxygen consumption rates when ambient $p\text{O}_2$ declines. However, these animals have to rely on anaerobic energy generation once $p\text{O}_{2,\text{crit}}$ is reached as well. Onset of anaerobic metabolism marks time-limited survival. The depletion of fermentable substrate (e.g., glycogen reserves) eventually results in catastrophic drops in cellular [ATP], which induces a cascade of events that lead to depolarization of membranes, Ca^{2+} entry into the cytoplasm and cell death (see Boutilier 2001 for review). Critical $p\text{O}_2$

values for heterotrophic metazoans are species specific and typically reached at an air saturation between 5 and 40 % (ca. 1–8 kPa; e.g., Grieshaber et al. 1994; Cochran and Burnett 1996; Zielinski et al. 2000; Nilsson et al. 2007; Mandic et al. 2009). However, effects of hypoxia on energy allocation and energy budgets (e.g., growth performance, reproductive effort) can usually be observed well before $pO_{2,crit}$ is reached and may be more important in defining ecological fitness (see Gray et al. 2002 for review).

Few benthic marine animals can survive prolonged periods of severe hypoxia or anoxia. Among the hypoxia tolerant, many annelids and mollusks are characterized by specific physiological adaptations that enable them to use limited energy stores more efficiently during anoxia. While the final cause of death is due to thermodynamic imbalances in cellular energy metabolism in both hypoxia-tolerant and hypoxia-intolerant species (see above), the former are adapted to downregulate cellular energy demand ('metabolic depression', Hochachka 1986; Guppy and Withers 1999; Guppy 2004) and to use anaerobic metabolic pathways that are more energy-yielding (Grieshaber et al. 1994; Hochachka and Lutz 2001) via down-regulation of cellular ATP and thus oxygen demand by a factor of 10–100 and increases in ATP yield by factors of 2–3 through the utilization of, for example, succinate and propionate as anaerobic end products, such species can survive anoxia on limited glycogen stores for several weeks (e.g., Dries and Theede 1974; Oeschger 1990; DeZwaan et al. 1991; Grieshaber et al. 1994; Boutilier 2001). In the deeper regions of Kiel Bay, for example, benthic communities are dominated by bivalves (*Astarte* spp. and *Arctica islandica*) that can survive weeks of anoxia by utilizing these key biochemical mechanisms: during a 50-day anoxic incubation, *A. islandica* was shown to accumulate high tissue concentrations of succinate (ca. 200 $\mu\text{mol g}^{-1}$ dry mass) while simultaneously reducing aerobic metabolic rate by more than 99 % (Oeschger 1990).

Hypercapnic hypoxia

As indicated above, the majority of laboratory experimental work on hypoxia tolerance of marine organisms has been carried out without the simultaneous manipulation of pCO_2 (see references discussed in Grieshaber et al. 1994; Gray et al. 2002; Levin et al. 2009). While it seems clear that hypoxia and hypercapnia both can impact animal energy budgets and thus might act additively or synergistically (see above), few studies have independently quantified hypoxia versus hypercapnia effects. Cochran and Burnett (1996) found that hypercapnia does not alter $pO_{2,crit}$ in two estuarine fish and a crustacean (shrimp) species. Similarly, Rosa and Seibel (2008) found

pO_2 and temperature, but not pCO_2 , to be the dominating factors determining metabolic rate reductions in a pelagic squid species that undergoes vertical migrations into oxygen-minimum zones. Other studies suggest that elevated pCO_2 and hypoxia may act synergistically in reducing oyster fitness by stimulating parasitism and decreasing defense mechanisms: Boyd and Burnett (1999) measured a decreased production of reactive oxygen species (ROS) in oyster blood cells (hemocytes). ROS are used by hemocytes in the defense against foreign cells, such as pathogenic bacteria. In addition, Dwyer and Burnett (1996) suggested that acidic conditions favor the growth of a protozoan parasite in oysters. Further work from Burnett's working group suggests an impairment of immune functions in estuarine organisms when hypoxia is combined with hypercapnia (e.g., Boyd and Burnett 1999; Mikulski et al. 2000). On the other hand, CO_2 also serves as an important adaptive signal during hypoxia: increases in the affinity of respiratory pigments (hemocyanins) toward oxygen in crustacean species are caused by elevated pCO_2 (Booth et al. 1982; Mangum and Burnett 1986). In fish, elevated pCO_2 elicits a ventilatory response that is thought to be mediated through branchial gas receptors (e.g., Gilmour 2001; Vulesevic et al. 2006). Finally, hypercapnia has also been suggested to aid in eliciting beneficial metabolic depression by altering extra- and/or intracellular acid-base status (see Guppy and Withers 1999).

Clearly, the magnitude of acidification that can currently be expected in hypoxic coastal areas (i.e., $pCO_2 > 1,700\text{--}3,200 \mu\text{atm}$, Fig. 5d, see also Feely et al. 2010: Pacific urban estuary, Cai et al. 2011: Gulf of Mexico, East China Sea) is greater than what is predicted for the next few 100 years due to ocean acidification in the oxygenated surface ocean (Caldeira and Wickett 2003, 2005). Linear increases in C_T due to respiration lead to exponential increases in pCO_2 , the main physiological effector for heterotrophic metazoans (Fig. 6). Wind-driven upwelling can bring hypoxic and hypercapnic waters from below pycnoclines in contact with productive shallow water ecosystems. Figure 4 illustrates wind-driven upwelling of low-pH water in Kiel Fjord, and Thomsen et al. (2010) showed that the general pattern of seasonal pH and pCO_2 found in Kiel Fjord surface water, although more variable, resembles that of the bottom waters at Boknis Eck Time Series Station (Figs. 2, 3). Maximum surface pCO_2 values around subtidal mussel beds in Kiel Fjord surpassed 2,300 μatm in September 2008, and average values in July and August (the main reproductive season for many calcifying invertebrates) averaged ca. 1,000 μatm (Thomsen et al. 2010). As the Baltic Sea is characterized by many seasonal and permanent hypoxic zones (HELCOM 2009) and wind-driven upwelling is common (e.g., Lehmann

et al. 2002; Myrberg and Andrejev 2003), it can be expected that a similar variability in carbonate system speciation is typical across the Baltic. Similar hypoxia-induced increases in $p\text{CO}_2$ will be characteristic for a multitude of coastal habitats worldwide (e.g., Diaz and Rosenberg 2008; see Cai et al. 2011 for a recent description of carbonate system speciation in East China Sea and Gulf of Mexico hypoxic zones).

Future ocean acidification and hypoxia

While the magnitude of acidification associated with low-oxygen conditions that can be attained under current conditions is already sizable, progressing invasion of anthropogenic CO_2 , that is, ocean acidification, will further elevate seawater $p\text{CO}_2$ in hypoxic regions in a nonlinear fashion (Fig. 5). This has recently also been pointed out by Brewer and Peltzer (2009) who estimated future changes in $p\text{CO}_2$ in oxygen-minimum zones that correspond to magnitudes of change simulated here (Fig. 3). Our model calculations for the $2\times \text{CO}_2$ scenario show that coastal hypoxic zones may easily encounter $p\text{CO}_2$ values of 3,400–4,500 μatm when most oxygen is consumed (Fig. 5d). Other habitats enriched in C_T already now, for example oceanic oxygen-minimum zones (Keeling et al. 2010; Paulmier et al. 2011), continental upwelling systems (Feely et al. 2008), estuaries (Borges et al. 2004; Feely et al. 2010; Waldbusser et al. 2011) and estuarine salt marsh systems (Wang and Cai 2004), will likely be impacted in a similar manner by future ocean acidification. It is important to stress that in oceanic oxygen-minimum zones, ocean acidification-enhanced $p\text{CO}_2$ will be encountered with a significant time delay, which is related to transit times of decades to a century (Karstensen et al. 2008), between last contact with the atmosphere and arrival in the OMZ. In coastal waters with seasonal hypoxia, this time delay is on the order of months to 1 year. Hence, such regions will feel the combined effect of ocean acidification and hypoxia almost instantaneously.

High $p\text{CO}_2$ values >3,400–4,500 μatm have been shown to negatively affect many heterotrophic metazoans in carbonate system perturbation experiments (see, e.g., Fabry et al. 2008; Ries et al. 2009; Dupont et al. 2010; Hofmann et al. 2010 for review). In order to maintain outward CO_2 diffusion gradients, animals will be challenged by severe acid–base disturbances due to proportionally higher extracellular $p\text{CO}_2$ values in hypoxic hypercapnic regions (Fig. 6). These acid–base disturbances may then significantly increase costs for cellular homeostasis, which would impact energy stores needed for survival under hypoxic or anoxic conditions (see above). Extremely low Ω_arag will be encountered in the future as well (Fig. 5c, f). This will represent an additional stressor, particularly for many

mollusks that dominate coastal benthic systems, as organic covers (periostraca) are typically thin or fractured in older specimens (Harper 1997; Thomsen et al. 2010). This will lead to enhanced rates of external shell dissolution. Future warming may exacerbate the problem by intensifying stratification, lowering saturating oxygen concentrations and elevating respiration rates, which individually and in combination could lead to a prolongation of hypoxic hypercapnic periods in seasonally hypoxic systems. This could then deplete the resistance capacity of even the most stress-tolerant species: during a severe hypercapnic hypoxic event in Kiel Bay in 1981, a 90 % decline in benthic biomass could be witnessed, with only few, highly anoxia-tolerant species surviving this period (the bivalves *Arctica islandica*, *Astarte* spp. and a few other species, Weigelt and Rumohr 1986). Additionally, it has been previously shown that elevated temperature results in reduced survival times under anoxia even in anoxia-tolerant bivalve species (Dries and Theede 1974). Clearly, the combined action of temperature, hypoxia and hypercapnia will constitute a severe challenge for benthic coastal ecosystems of the future (e.g., Pörtner et al. 2005).

Conclusions and implications for future research

Our basic calculations illustrate that hypoxic coastal areas are already characterized by $p\text{CO}_2$ values that will probably not be reached by ocean acidification in the surface ocean in the next few 100 years (i.e., depending on salinity, >1,700–3,200 μatm), with the potential for a 50–100 % increase within this century. Particularly high $p\text{CO}_2$ values and very low Ω_arag can be expected in highly productive estuarine habitats. Yet, little attention has been paid to the magnitudes and rates of change in $p\text{CO}_2$ in these systems. Studies that investigate the effects of hypoxia on benthic ecosystems and species should focus energy and resources toward:

1. Studying the carbonate system speciation in hypoxic habitats in relation to $[\text{O}_2]$ and temperature variation.
2. Studying potential upwelling events and seasonal dynamics of variability in carbonate system speciation.
3. Studying species and community responses under simultaneous application of realistic $[\text{O}_2]$, $[\text{CO}_2]$ and temperature combinations. This recommendation refers to both rate and magnitude of stress application. CO_2 can act both as a stressor and as an adaptive modulator of physiological function.

In ocean acidification research, it is often recommended to primarily utilize $p\text{CO}_2$ values between current and up to ca 1,000 μatm in experimental CO_2 perturbation experiments (based on IPCC scenarios, e.g., IPCC 2007).

However, respiration processes can significantly elevate $p\text{CO}_2$ in coastal waters, and upwelling of severely hypercapnic and hypoxic water may further elevate $p\text{CO}_2$. Additional CO_2 through ocean acidification can produce much higher $p\text{CO}_2$ values than predicted by the global surface ocean model predictions (e.g., Caldeira and Wickett 2005). Thus, experimental $p\text{CO}_2$ levels should be assigned in a case-sensitive manner after the specific carbonate system variability of the habitat in question has been studied. In a next step, future developments can be roughly estimated using standard carbonate system speciation software and applied to perturbation experiments.

It also should be kept in mind that hypoxic hypercapnic systems and upwelling systems as described here can serve as interesting model regions where responses of adapted marine ecosystems toward high levels of acidification can already be studied today (c.f. Thomsen et al. 2010). These systems may tell us much about animal tolerance limits in response to an altered carbonate system speciation. Whether these ecosystems also can cope with the high levels of acidification estimated here to occur within this century remains to be established.

Acknowledgments F. Melzner and A. Körtzinger were supported by the DFG Excellence Cluster project ‘Future Ocean’ CP0816: ‘Carbonate system variability in Kiel Bay’; F. Melzner, M.A. Gutowska, W. Koeve and A. Oschlies are supported by the German National Project ‘Biological impacts of ocean acidification (BIO-ACID)’, funded by BMBF. The Boknis Eck Time Series Station (BE) is operated by the Chemical Oceanography department at IFM-GEOMAR: www.ifm-geomar.de/index.php?id=bokniseck. The authors would like to thank the crew of FK ‘Littorina’ and all researchers and students involved in the BE data collection efforts between 1986 and 1995, especially G. Behrends, H. Giesenhausen, U. Horstmann and H. Johansen. Between 1985 and 1992, BE was funded by BMFT project 03F0547A (‘Biologisches Monitoring der Ostsee’) and between 1993 and 1996 by BMFT project 03F0166A (‘Kieler Bucht Projekt’).

References

- Beldowski J, Löffler A, Schneider B, Joensuu L (2010) Distribution and biogeochemical control of total CO_2 and total alkalinity in the Baltic Sea. *J Mar Syst* 81:252–259
- Booth CE, McMahon BR, Pinder AW (1982) Oxygen uptake and the potentiating effects of increased hemolymph lactate on oxygen transport during exercise in the blue crab, *Callinectes sapidus*. *J Comp Physiol* 148:111–1216
- Borges AV, Delille B, Schiettecatte LS, Gazeau F, Abril G, Frankignoulle M (2004) Gas transfer velocities of CO_2 in three European estuaries (Randers Fjord, Scheldt, and Thames). *Limnol Oceanogr* 49:1630–1641
- Boutillier RG (2001) Mechanisms of cell survival in hypoxia and hypothermia. *J Exp Biol* 204:3171–3181
- Boyd JN, Burnett LE (1999) Reactive oxygen intermediate production by oyster hemocytes exposed to hypoxia. *J Exp Biol* 202:3135–3143
- Brewer PG (1997) Ocean chemistry of the fossil fuel CO_2 signal: the haline signal of “business as usual”. *Geophys Res Lett* 24:1367–1369
- Brewer PG, Peltzer ET (2009) Limits to marine life. *Science* 324:347–348
- Burnett LE (1997) The challenges of living in hypoxic and hypercapnic aquatic environments. *Am Zool* 37:633–640
- Cai WJ, Hu X, Huang WJ, Murrell MC, Lehrter JC, Lohrenz SE, Chou WC, Zhai W, Hollibaugh JT, Wang Y, Zhao P, Guo X, Gundersen K, Dai M, Gong GC (2011) Acidification of subsurface coastal waters enhanced by eutrophication. *Nat Geosci* 4:766–770
- Caldeira K, Wickett ME (2003) Anthropogenic carbon and ocean pH. *Nature* 425:365
- Caldeira K, Wickett ME (2005) Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean. *J Geophys Res* 110:C09S04. doi:10.1029/2004JC002671
- Cao L, Caldeira K (2008) Atmospheric CO_2 stabilization and ocean acidification. *Geophys Res Lett* 35:L19609. doi:10.1029/2008GL035072
- Cochran RE, Burnett LE (1996) Respiratory responses of the salt marsh animals, *Fundulus heteroclitus*, *Leiostomus xanthurus*, and *Palaemonetes pugio* to environmental hypoxia and hypercapnia and to the organophosphate pesticide, azinphosmethyl. *J Exp Mar Biol Ecol* 195:125–144
- Deigweier K, Hirse T, Bock C, Lucassen M, Pörtner HO (2009) Hypercapnia induced shifts in gill energy budgets of Antarctic notothenioids. *J Comp Physiol B* 180:347–359
- DeZwaan A, Cortesi P, Van den Thillart G, Roos J, Storey KB (1991) Differential sensitivities to hypoxia by 2 anoxia-tolerant marine mollusks—a biochemical analysis. *Mar Biol* 111:343–351
- Diaz RJ, Rosenberg R (1995) Marine benthic hypoxia: a review of its ecological effects and the behavioural responses of benthic macrofauna. *Oceanogr Mar Biol* 33(33):245–303
- Diaz RJ, Rosenberg R (2008) Spreading dead zones and consequences for marine ecosystems. *Science* 321:926–929
- Dickson AG (1990) Standard potential of the reaction: $\text{AgCl}_s + 1/2 \text{H}_2\text{g} = \text{Ag}_s + \text{HCl}_{\text{aq}}$, and the standard acidity constant of the ion HSO_4^- in synthetic seawater from 273.15 to 318.15 K. *J Chem Thermodyn* 22:113–127
- Dickson AG, Sabine CL, Christian JR (2007) Guide to best practices for ocean CO_2 measurements, vol 3. PICES Special Publications, p 191
- Dixson DL, Munday PL, Jones GP (2010) Ocean acidification disrupts the innate ability of fish to detect predator olfactory cues. *Ecol Lett* 13:68–75
- Dries RR, Theede H (1974) Resistance to oxygen deficiency in marine bottom invertebrates of the Western Baltic Sea. *Mar Biol* 25:327–333
- Dupont S, Ortega-Martinez O, Thorndyke M (2010) Impact of near-future ocean acidification on echinoderms. *Ecotoxicology* 19:449–463
- Dwyer JJ, Burnett LE (1996) Acid-base status of the oyster *Crassostrea virginica* in response to air exposure and to infections by *Perkinsus marinus*. *Biol Bull* 190:139–147
- Ellington WR (1993) Studies of intracellular pH regulation in cardiac myocytes from the marine bivalve mollusk, *Mercenaria campechiensis*. *Biol Bull* 184:209–215
- Fabry VJ, Seibel BA, Feely RA, Orr JC (2008) Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES J Mar Sci* 65:414–432
- Farrell AP, Richards JG (2009) Defining hypoxia: an integrative synthesis of the responses of fish to hypoxia. In: Richards JG, Farrell AP, Brauner CJ (eds) Hypoxia. Academic Press, London, pp 487–503. doi:10.1016/S1546-5098(08)00011-3
- Feely RA, Sabine CL, Hernandez-Ayon JM, Ianson D, Hales B (2008) Evidence for upwelling of corrosive “acidified” water onto the continental shelf. *Science* 320:1490–1492

- Feely RA, Alin SR, Newton J, Sabine CL, Warner M, Devol A, Krembs C, Maloy C (2010) The combined effects of ocean acidification, mixing, and respiration on pH and carbonate saturation in an urbanized estuary. *Estuar Coast Shelf S* 88:442–449
- Frankignoulle M, Bourge I, Wollast R (1996) Atmospheric CO₂ fluxes in a highly polluted estuary (the Scheldt). *Limnol Oceanogr* 41:365–369
- Gilmour KM (2001) The CO₂/pH ventilatory drive in fish. *Comp Biochem Physiol A* 130:219–240
- Grashoff K, Ehrhardt M, Kremling K (eds) (1983) *Methods of seawater analysis*. Weinheim/Deerfield Beach, Florida, p 419. ISBN:3-527-2599-8 (Weinheim) 0-89573-7 (Deerfield Beach)
- Gray JS, Wu RSS, Or YY (2002) Effects of hypoxia and organic enrichment on the coastal marine environment. *Mar Ecol Prog Ser* 238:249–279
- Grieshaber M, Hardewig I, Kreutzer U, Pörtner HO (1994) Physiological and metabolic responses to hypoxia in invertebrates. *Rev Physiol Biochem Pharmacol* 125:43–147
- Guppy M (2004) The biochemistry of metabolic depression: a history of perceptions. *Comp Biochem Physiol B: Biochem Mol Biol* 139:435–442
- Guppy M, Withers P (1999) Metabolic depression in animals: physiological perspectives and biochemical generalizations. *Biol Rev* 74:1–40
- Gutowska MA, Melzner F, Pörtner HO, Meier S (2010) Cuttlebone calcification increases during exposure to elevated seawater pCO₂ in the cephalopod *Sepia officinalis*. *Mar Biol* 157:1653–1663
- Hansen HP, Giesenhausen HC, Behrends G (1999) Seasonal and long-term control of bottom-water oxygen deficiency in a stratified shallow-water coastal system. *ICES J Mar Sci* 56:65–71
- Harper EM (1997) The molluscan periostracum: an important constraint in bivalve evolution. *Palaeontology* 40:71–97
- HELCOM (2009) Eutrophication in the Baltic Sea—an integrated thematic assessment of the effects of nutrient enrichment and eutrophication in the Baltic Sea region. *Baltic Sea Environmental Proceedings No. 115B*, ISSN 0357-2994
- Hochachka PW (1986) Defense strategies against hypoxia and hypothermia. *Science* 231:234–241
- Hochachka PW, Lutz PL (2001) Mechanism, origin, and evolution of anoxia tolerance in animals. *Comp Biochem Physiol B* 130:435–459
- Hofmann GE, Barry JP, Edmunds PJ, Gates RD, Hutchins DA, Klinger T, Sewell MA (2010) The effect of ocean acidification on calcifying organisms in marine ecosystems: an organism-to-ecosystem perspective. *Annu Rev Ecol Evol Syst* 41:127–147. doi:10.1146/annurev.ecolsys.110308.120227
- Hofmann AF, Peltzer ET, Walz PM, Brewer PG (2011) Hypoxia by degrees: establishing definitions for a changing ocean. *Deep-Sea Res I* 58:1212–1226
- IPCC (2007) *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge
- Karstensen J, Stramma L, Visbeck M (2008) Oxygen minimum zones in the eastern tropical Atlantic and Pacific oceans. *Prog Oceanogr* 77:331–350
- Keeling R, Körtzinger A, Gruber N (2010) Ocean deoxygenation in a warming world. *Ann Rev Mar Sci* 2:463–493
- Kleypas JA, Buddemeier RW, Archer D, Gattuso JP, Langdon C, Opdyke BN (1999) Geochemical consequences of increased atmospheric carbon dioxide on coral reefs. *Science* 284:118–120
- Körtzinger A, Hedges JJ, Quay PD (2001) Redfield ratios revisited: removing the biasing effect of anthropogenic CO₂. *Limnol Oceanogr* 46:964–970
- Lehmann A, Krauss W, Hinrichsen HH (2002) Effects of remote and local atmospheric forcing on circulation and upwelling in the Baltic Sea. *Tellus Ser Dyn Meteorol Oceanogr* 54:299–316
- Levin LA, Ekau W, Gooday AJ et al (2009) Effects of natural and human-induced hypoxia on coastal benthos. *Biogeosciences* 6:2063–2098
- Lewis E, Wallace DWR (1998) Program developed for CO₂ system calculations. Oak Ridge, Oak Ridge National Laboratory ORNL/CDIAC-105
- Lischka S, Büdenbender J, Boxhammer T, Riebesell U (2011) Impact of ocean acidification and elevated temperatures on early juveniles of the polar shelled pteropod *Limacina helicina*: mortality, shell degradation, and shell growth. *Biogeosciences* 8:919–932
- Mandic M, Sloman KA, Richards JG (2009) Escaping to the surface: a phylogenetically independent analysis of hypoxia-induced respiratory behaviors in sculpins. *Physiol Biochem Zool* 82:730–738
- Mangum CP, Burnett LE (1986) The CO₂ sensitivity of the hemocyanins and its relationship to Cl[−] sensitivity. *Biol Bull* 171:248–263
- Melzner F, Gutowska MA, Langenbuch M et al (2009) Physiological basis for high CO₂ tolerance in marine ectothermic animals: pre-adaptation through lifestyle and ontogeny? *Biogeosciences* 6:2313–2331
- Melzner F, Stange P, Trübenbach K, Thomsen J, Casties I, Panknin U, Gorb SN, Gutowska MA (2011) Food supply and seawater pCO₂ impact calcification and internal shell dissolution in the blue mussel *Mytilus edulis*. *PLoS ONE* 6(9):e24223. doi:10.1371/journal.pone.0024223
- Mikulski CM, Burnett LE, Burnett KG (2000) The effects of hypercapnic hypoxia on the survival of shrimp challenged with *Vibrio parahaemolyticus*. *J Shellfish Res* 19:301–311
- Mintrop L, Perez FF, Gonzalez-Davila M, Santana-Casiano MJ, Körtzinger A (2000) Alkalinity determination by potentiometry: intercalibration using three different methods. *Cienc Mar* 26:23–37
- Munday PL, Dixon DL, Donelson JM, Jones GP, Pratchett MS, Devitsina GV, Doving KB (2009) Ocean acidification impairs olfactory discrimination and homing ability of a marine fish. *Proc Nat Acad Sci USA* 106:1848–1852
- Myrberg K, Andrejev O (2003) Main upwelling regions in the Baltic Sea—a statistical analysis based on three-dimensional modeling. *Boreal Environ Res* 8:97–112
- Nilsson GE, Hobbs JPA, Ostlund-Nilsson S (2007) Tribute to P. L. Lutz: respiratory ecophysiology of coral-reef teleosts. *J Exp Biol* 210:1673–1686
- Nilsson GE, Dixon DL, Domenici P, McCormick MI, Christina Sørensen C, Watson SA, Munday PL (2012) Near-future carbon dioxide levels alter fish behaviour by interfering with neurotransmitter function. *Nat Clim Chang* 2:201–204. doi:10.1038/nclimate1352
- Oeschger R (1990) Long-term anaerobiosis in sublittoral marine invertebrates from the Western Baltic Sea—*Halicryptus spinulosus* (Priapulida), *Astarte borealis* and *Arctica islandica* (Bivalvia). *Mar Ecol Progr Ser* 59:133–143
- Paulmier A, Ruiz-Pino D, Garçon V (2011) CO₂ maximum in the oxygen minimum zone (OMZ). *Biogeosciences* 8:239–252
- Pörtner HO, Grieshaber MK (1993) Critical PO₂(s) in oxyconforming and oxyregulating animals gas exchange, metabolic rate and the mode of energy production. In: Bicudo JEPW (ed) *The vertebrate gas transport cascade adaptations to environment and mode of life*. CRC Press, Boca Raton, pp 330–357
- Pörtner HO, Langenbuch M, Michaelidis B (2005) Synergistic effects of temperature extremes, hypoxia, and increases in CO₂ on marine animals: from earth history to global change. *J Geophys Res Oceans* 110:C09S10

- Rabalais NN, Turner RE, Wiseman WJ (2002) Gulf of Mexico hypoxia, aka “The dead zone”. *Annu Rev Ecol Syst* 33:235–263
- Richards JG (2009) Metabolic and molecular responses of fish to hypoxia. In: Richards JG, Farrell AP, Brauner CJ (eds) *Hypoxia*. Academic, London, pp 443–485. doi:[10.1016/S1546-5098\(08\)00010-1](https://doi.org/10.1016/S1546-5098(08)00010-1)
- Ries JB, Cohen AL, McCorkle DC (2009) Marine calcifiers exhibit mixed responses to CO₂-induced ocean acidification. *Geology* 37:1131–1134
- Rosa R, Seibel BA (2008) Synergistic effects of climate-related variables suggest future physiological impairment in a top oceanic predator. *Proc Nat Acad Sci USA* 105:20776–20780
- Roy RN, Roy LN, Vogel KM, Portermore C, Pearson T, Good CE, Millero FJ, Campbell DM (1993) The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45 °C. *Mar Chem* 44:249–267
- Seibel BA (2011) Critical oxygen levels and metabolic suppression in oceanic oxygen minimum zones. *J Exp Biol* 204:326–336
- Seibel BA, Walsh PJ (2003) Biological impacts of deep-sea carbon dioxide injection inferred from indices of physiological performance. *J Exp Biol* 206:641–650
- Stump M, Wren J, Melzner F, Thorndyke MC, Dupont S (2011) CO₂ induced seawater acidification impacts sea urchin larval development I: elevated metabolic rates decrease scope for growth and induce developmental delay. *Comp Biochem Phys A* 160:331–340
- Thomsen J, Melzner F (2010) Moderate seawater acidification does not elicit long-term metabolic depression in the blue mussel *Mytilus edulis*. *Mar Biol* 157:2667–2676
- Thomsen J, Gutowska MA, Saphörster J, Heinemann A, Trübenbach K, Fietzke J, Hiebenthal C, Eisenhauer A, Körtzinger A, Wahl M, Melzner F (2010) Calcifying invertebrates succeed in a naturally CO₂-rich coastal habitat but are threatened by high levels of future acidification. *Biogeosciences* 7:3879–3891
- Tomanek L, Zuzow MJ, Ivanina AV, Beniash E, Sokolova IM (2011) Proteomic response to elevated PCO₂ level in eastern oysters, *Crassostrea virginica*: evidence for oxidative stress. *J Exp Biol* 214:1836–1844
- Truchot JP, Duhameljoue A (1980) Oxygen and carbon dioxide in the marine intertidal environment—diurnal and tidal changes in rockpools. *Respir Physiol* 39:241–254
- Tunnicliffe V, Davies KTA, Butterfield DA, Embley RW, Rose JM, Chadwick WW Jr (2009) Survival of mussels in extremely acidic waters on a submarine volcano. *Nat Geosci* 2:344–348
- Vaquier-Sunyer R, Duarte CM (2008) Thresholds of hypoxia for marine biodiversity. *Proc Nat Acad Sci USA* 105:15452–15457
- Vulesevic B, McNeill B, Perry SF (2006) Chemoreceptor plasticity and respiratory acclimation in the zebrafish *Danio rerio*. *J Exp Biol* 209:1261–1273
- Waldbusser GG, Voight EP, Bergschneider H, Green MA, Newell RIE (2011) Biocalcification in the Eastern Oyster (*Crassostrea virginica*) in relation to long-term trends in Chesapeake Bay pH. *Estuar Coast* 34:221–231
- Wang Z, Cai WJ (2004) Carbon dioxide degassing and inorganic carbon export from a marsh dominated estuary (the Duplin River): a marsh CO₂ pump. *Limnol Oceanogr* 49:341–352
- Weigelt M, Rumohr H (1986) Effects of wide-range oxygen depletion on benthic fauna and demersal fish in Kiel bay 1981–1983. *Meeresforschung Rep Mar Res* 31:124–136
- Weiner S, Addadi L (2011) Crystallization pathways in biomineralization. *Annu Rev Mater Res* 41:21–40
- Widdicombe S, Spicer JI (2008) Predicting the impact of ocean acidification on benthic biodiversity: what can animal physiology tell us? *J Exp Mar Biol Ecol* 366:187–197
- Wood HL, Spicer JI, Widdicombe S (2008) Ocean acidification may increase calcification rates, but at a cost. *Proc R Soc B Biol Sci* 275:1767–1773
- Yates KK, Dufore C, Smiley N, Jackson C, Halley RB (2007) Diurnal variation of oxygen and carbonate system parameters in Tampa Bay and Florida Bay. *Mar Chem* 104:110–124. doi:[10.1016/j.marchem.2006.12.008](https://doi.org/10.1016/j.marchem.2006.12.008)
- Zange J, Grieshaber MK, Jans AW (1990) The regulation of intracellular pH estimated by ³¹P-NMR spectroscopy in the anterior byssus retractor muscle of *Mytilus edulis* L. *J Exp Biol* 150:95–109
- Zielinski S, Lee PG, Pörtner HO (2000) Metabolic performance of the squid *Lolliguncula brevis* (Cephalopoda) during hypoxia: an analysis of the critical PO₂. *J Exp Mar Biol Ecol* 243:241–259